

How to Mix Two Vials of 5,000 IU Novarel hCG to Give One Injection of 10,000 IU Novarel hCG

STEP 1: Remove the plastic caps from two vials of 5,000 IU Novarel hCG and one (1) vial of bacteriostatic water (also known as diluent).

STEP 2: Swab the top of each of the three vials with alcohol wipe.

STEP 3: Using a 3mL syringe with 1 ½" needle, withdraw 1mL of the diluent.

STEP 4: Inject that 1mL of diluent into the first vial of 5,000 IU hCG to reconstitute the powder and immediately withdraw the entire amount back into the syringe.

STEP 5: Inject the contents of that syringe into the second vial of 5,000 IU hCG. Make sure the reconstituted powder is completely dissolved. Withdraw the entire amount back into the syringe.

You have now mixed two vials of 5,000 IU Novarel hCG into one single 10,000 IU injection.

Replace mixing needle with a new 1 ½" needle for IM injection, or ½" needle for SQ injection.

Please view our website <https://www.vfp-pharmacygroup.com/education/medication-instructional-videos/> for IM or SQ instructional videos.

If your dose is less than 10,000 IU, please obtain specific mixing instructions from your clinic, or contact your VFP Pharmacy Group Location at: <https://www.vfp-pharmacygroup.com/contact-us/#our-locations>

.